

# 668P: First-in-Human Phase I Dose Escalation Study of ALG.APV-527, a 5T4 Tumor Antigen-Conditional 4-1BB Bispecific Antibody, in Patients with Advanced Solid Tumors, Demonstrates Positive Safety, Signals of Biological Activity and Patients with Lasting Stable Disease




Thomas Marron<sup>1</sup>, Seth Rosen<sup>2</sup>, Douglas Orr<sup>3</sup>, John Powderly II<sup>4</sup>, Jennifer Segar<sup>5</sup>, Noor Khaskhely<sup>6</sup>, Lynn Bonham<sup>6</sup>, Michelle Nelson<sup>6</sup>, Caroline Taromino<sup>6</sup>, Daphne Taylor<sup>6</sup>, John Kumer<sup>6</sup>, Peter Ellmark<sup>7,8</sup>, Sara Fritzell<sup>7</sup>, Sumeet Ambarkhane<sup>7</sup>, Dirk Huebner<sup>6</sup>, Asrar Alahmadi<sup>9</sup>




1.Tisch Cancer Center, Icahn School of Medicine at Mount Sinai, New York, NY, USA; 2.Hematology Oncology Associates of the Treasure Coast, Port Saint Lucie, FL, USA; 3.Mary Crowley Cancer Research, Dallas, TX, USA; 4.Carolina BioOncology Institute, Huntersville, NC, USA; 5.University of Arizona Comprehensive Cancer Center, Tucson, AZ, USA; 6.Aptevo Therapeutics, Seattle, WA, USA; 7.Alligator Bioscience AB, Lund, Sweden; 8.Department of Immunotechnology, Lund University, Lund, Sweden; 9.Ohio State University Comprehensive Cancer Center, Columbus, OH, USA

## About ALG.APV-527

- ALG.APV-527 is a bispecific therapeutic containing binding domains targeting the co-stimulatory receptor 4-1BB and the oncofetal antigen 5T4, expressed on multiple solid tumor types. These are linked to an effector-null Ig Fc domain, providing an antibody-like *in vivo* half-life
- The scFvs originate from the Alligator Gold<sup>®</sup> human scFv library (Alligator Bioscience) and optimized for use in the bispecific ADAPTIR<sup>™</sup> format (Aptevo Therapeutics)
- ALG.APV-527 features targeted T cell stimulation, optimized stability, good manufacturing properties with potential for better risk-benefit in humans than monospecific 4-1BB antibodies



## ALG.APV-527 Mode of Action



ALG.APV-527 directs the stimulation of CD8<sup>+</sup> T and NK cells by 5T4<sup>+</sup> tumors and is designed to minimize the toxicity observed with other 4-1BB therapeutics

## Study Design

The Phase I study is a first-in-human, open-label, multicenter trial consisting of up to six cohorts (0.1-15 mg/kg) with a 3+3 dose escalation of ALG.APV-527 monotherapy, administered IV Q2W, in adult patients with advanced solid tumors. Eligibility is limited to patients with tumor types identified as likely to express 5T4 antigen.

Clinical Trials Number: NCT05934539

### Key Objectives

- Characterize **safety & tolerability profile** of ALG.APV-527
- Identify **MTD** and/or **RP2D**
- Characterize **PK profile** after single and repeated IV administration
- Assess potential **immunogenicity** and **PD** effects
- Obtain a preliminary assessment of **anti-tumor activity**

## Baseline Characteristics

Dose mg/kg Total n=18	Median age (range)	Gender (%)		ECOG PS (%)		Tumor Types
		F	M	0	1	
Cohort 1 0.1 (n=4)	75.5 (71-82)	2 (50)	2 (50)	0	4 (100)	Breast, Colorectal (2) Esophagus
Cohort 2 0.5 (n=3)	58.0 (56-74)	1 (33)	2 (67)	0	3 (100)	Breast, Pancreatic (2)
Cohort 3 2.0 (n=6)	56.5 (39-72)	5 (83)	1 (17)	2 (33)	4 (67)	H&N SCC, Pancreatic Colorectal (4)
Cohort 4 6.0 (n=3)	47.0 (44-60)	2 (67)	1 (33)	1 (33)	2 (67)	NSCLC, Colorectal (2)
Cohort 5 12.0 (n=2)	56.0 (51-61)	2 (100)	0	0	2 (100)	Renal CC, NSCLC

Prior Anti-Cancer Therapy	n (%); median (range)	Prior Systemic Therapy	n (%)
Surgery	8 (44); 1 (1-3)	Chemotherapy & other antineoplastic therapies	18 (100)
Radiotherapy	12 (67); 1 (1-3)	Immune and antibody therapies*	13 (72)
Systemic Therapy	18 (100); 6 (2-8)		

n= number of patients; \*bevacizumab (8), pembrolizumab (3), nivolumab (2), ipilimumab, IL-2, cetuximab, panitumumab, durvalumab, cemiplimab


## Treatment-Related AEs

>1 Patient; 18 total patients	n (%) E	Serious Treatment-Related AEs	n (%) E
Any TRAE	16 (89) 87	Any Serious TRAE	4 (22) 5
Fatigue	4 (22) 6	Febrile neutropenia*	1 (6) 1
Diarrhea	3 (17) 4	Ulcerative colitis	1 (6) 1
Infusion-related reaction	3 (17) 6	Hemorrhagic diarrhea	1 (6) 1
Nausea	2 (11) 2	GI hemorrhage	1 (6) 1
ALT increase	2 (11) 2	Nausea	1 (6) 1
Anemia	2 (11) 4		
Neutrophil count decrease	2 (11) 2		
WBC count decrease	2 (11) 3		
Myalgia	2 (11) 2		
Pruritus	2 (11) 3		


Overall, ALG.APV-527 was safe and well tolerated.

\*Case of Gr4 febrile neutropenia, classified as a DLT, and resolved following Filgrastim treatment.  
n= number of patients, E= number of events

## Favorable PK profile and increase in soluble 4-1BB in blood following treatment



## Duration on Study



Fifteen of 18 patients were evaluable at data cutoff (26 July 2024). Nine patients had a best overall response of SD. Cohort 2 breast cancer patient increased dosing from 0.5 to 2.0, then to 6.0 mg/kg.

SD= stable disease, PD= progressive disease \*Patients were allowed to stay on study with PD events if agreed upon with the treating physician

## Summary and Conclusions

ALG.APV-527 demonstrates good tolerability, safety, and biological activity. Serum concentration of ALG.APV-527 was consistent with the administered dose. Based on RECIST 1.1 and iRECIST, nine of 15 evaluable patients (60%) had a best overall response of SD, with the longest SD duration in a breast cancer patient >11 months. One colon cancer patient with sustained SD is still on study. Dose escalation is ongoing. An MTD has yet to be determined.